arXiv:1410.8511v1 [math.CV] 30 Oct 2014

SUBORDINATION PRINCIPLE FOR PROPER FUNCTIONS

JAN WIEGERINCK AND DJIRE IBRAHIM K.

Abstract. Our gaol is to generalize Littlewood's Subordination Theorem to the situations where the functions are not globally subordinate. On using our result we establish a relation between the moduli of the zeros of the subordinate function and the moduli of the zeros of the superordinate function. This fact has a consequence on the envelope of Lelong functional. At the end we give some coefficient inequalities of the subordinate and superordinate functions.

1. Introduction

Let f and g be two holomorphic functions, with $f(0) = g(0)$. Suppose that, f is bijective and $g(\mathbb{D}) \subset f(\mathbb{D})$. Then $\omega(z) = f^{-1}(g(z))$ is analytic in $\mathbb{D}$, $\omega(0) = 0$, $|\omega(z)| < 1$ and $g(z) = f \circ \omega(z)$. In general, an analytic function g is said to be subordinate to an analytic function f if $g(z) = f(\omega(z))$, $|z| < 1$, for some analytic function ω with $|\omega(z)| \leq |z|$. The superordinate function f need not be univalent. Littlewood's Subordination principle states that if f and g are analytic in the unit disk and if g is subordinate to f, then, for $0 < p < \infty$, $0 < r < 1$ we have:

$$\int_0^{2\pi} |g(re^{i\theta})|^p d\theta \leq \int_0^{2\pi} |f(re^{i\theta})|^p d\theta.$$

Under its general form we have for any subharmonic function u in some neighborhood of $f(\mathbb{D})$:

$$\int_0^{2\pi} u \circ g(re^{i\theta}) d\theta \leq \int_0^{2\pi} u \circ f(re^{i\theta}) d\theta.$$

For more detail see $[1, chap.6]$.

2. Generalization

We start by establishing some notation. For $r > 0$, let $D_r = \{z \in \mathbb{C}, |z| < r\}$ and $\mathbb{D} = D_1$. For $X \subset \mathbb{C}$ open, $SH(X)$ denotes the set of all subharmonic functions on X and $\mathscr{O}(D_r, X)$ the set of all holomorphic functions from D_r to X. An element $f \in \mathscr{O}(\mathbb{D}, X)$ is sometimes called an analytic disc in X of center $f(0)$. The following theorem generalizes Littlewood's Subordination principle to the situations where g is not subordinate to f.

Theorem 1. *Let X be an open subset in $\mathbb{C}$, and $f \in \mathscr{O}(\mathbb{D}, X) \cap C(\overline{\mathbb{D}}, X)$ proper, then for all $u \in SH(X)$ such that $|u \circ f|$ is bounded on $\mathbb{T}$, for all $g \in \mathscr{O}(\mathbb{D}, X)$ with $g(0) = f(0)$ and $g(\mathbb{D}) \subset f(\mathbb{D})$ one has*

$$\int_0^{2\pi} u \circ g(re^{i\theta}) d\theta \leq \int_0^{2\pi} u \circ f(e^{i\theta}) d\theta \text{ for all } r \in]0, 1[.$$

Key words and phrases. Analytic disc, subordination, subharmonic, Lelong functional.

Before proving this theorem we will recall a result concerning proper holomorphic functions.

Proposition 2. *Let $f \in \mathscr{O}(\mathbb{D}, \mathbb{C})$ proper, $w \in \Omega = f(\mathbb{D})$, then the set $f^{-1}(w)$ is finite and $\mathbb{D} \cap f'^{-1}(0)$ is countable.*

Under the assumptions in Theorem 1 we will use Perron method for the Dirichlet Problem on the closed unit disk $\overline{\mathbb{D}}$ to define a superharmonic function V on the open set $f(\mathbb{D})$ which is greater than u on $f(\mathbb{D})$ and we use the properties of superharmonic functions to infer the inequality in Theorem 1.

Proof of Theorem 1. Let $\epsilon > 0$, g , f and u be as in Theorem 1. As u is upper semicontinuous, then there is a sequence of continuous functions $(\psi_j)_j \subset C(X)$ which decreases to u. Set $M = \max\{\psi_1 \circ f(t), t \in \mathbb{T}\}$. As $|u \circ f(t)|$ is bounded on $\mathbb{T}$, then the function $[0, 2\pi] \ni \theta \to g(\theta) = \max\{|u \circ f(e^{i\theta})|, |M|\}$ is integrable on $[0, 2\pi]$. Notice that $|\psi_j \circ f(e^{i\theta})| \leq g(\theta)$ for all $\theta \in [0, 2\pi]$ and $j > 0$. Then by Lebesgue dominated convergence theorem there is j_0 such that:

$$\int_0^{2\pi} \psi_j \circ f(e^{i\theta}) d\theta < \int_0^{2\pi} u \circ f(e^{i\theta}) d\theta + \epsilon$$

for any $j > j_0$. We set $\psi = \psi_{j_0+1}$. Then $\int_0^{2\pi} \psi \circ f(e^{i\theta}) d\theta < \int_0^{2\pi} u \circ f(e^{i\theta}) d\theta + \epsilon$. We will define a superharmonic function V on $f(\mathbb{D})$ such that $u \leq V$ on $f(\mathbb{D})$. Set

$$v(t) = \sup\{w(t), w \in SH(\mathbb{D}), w^*(t) \leq \psi \circ f(t), t \in \mathbb{T}\},$$

then, v is harmonic on $\mathbb{D}$ and $v(t) = \psi \circ f(t)$ for $t \in \mathbb{T}$ see $[3, corol.4.1.8]$. Set $\Omega = f(\mathbb{D})$ and define $v_1 : \Omega \longrightarrow \mathbb{R}$ by

$$v_1(x) = \max\{-v(t), t \in f^{-1}(x)\}$$

where $x \in \Omega$ see [2]. By Proposition 2, $f^{-1}(x)$ is finite so v_1 makes sense. We will prove that v_1 is subharmonic on Ω. Let $x \in \Omega \setminus f(\mathbb{D} \cap f'^{-1}(0))$, then, by Local Inversion Theorem there exist a number $k \in \mathbb{N}$, a neighborhood W of x , disjoint neighbourhoods W_1 ,..., W_k of $\{t_1, ..., t_k\} = f^{-1}(x)$ and holomorphic functions $g_i : W \to W_i$ such that $f \circ g_i(y) = y$ for all $y \in W$. Then for all $y \in W$ we have

$$v_1(y) = \max\{-v \circ g_i(y), i = 1, .., k\}.$$

Hence v_1 is subharmonic on W, this for all $x \in \Omega \setminus f(\mathbb{D} \cap f'^{-1}(0))$. As subharmonicity is a local property then v_1 is subharmonic on $\Omega \setminus f(\mathbb{D} \cap f'^{-1}(0))$. As v_1 is bounded and $f(\mathbb{D} \cap f'^{-1}(0))$ is polar see $[3, corol.3.2.5]$, then by Removable Singularity Theorem, v_1 can be extended to a subharmonic function on Ω. One can find a similar work in $[4, page73]$.

We define a superharmonic function V on Ω on setting $V = -v_1$. Notice that

$$V(x) = \min\{v(t), t \in f^{-1}(x)\} \text{ and that } u \leq V \text{ on } \Omega.$$

We also have $u \circ f(t) \leq V \circ f(t) \leq v(t)$ for any $t \in \mathbb{D}$. As V is superharmonic, v is harmonic and $g(\mathbb{D}) \subset \Omega$, then we have

$$\frac{1}{2\pi} \int_0^{2\pi} u \circ g(re^{i\theta}) d\theta \leq \frac{1}{2\pi} \int_0^{2\pi} V \circ g(re^{i\theta}) d\theta \leq V \circ g(0) \leq v(0) = \frac{1}{2\pi} \int_0^{2\pi} \psi \circ f(e^{i\theta}) d\theta.$$

Hence by the choice of ψ we get

$$\int_0^{2\pi} u\circ g(re^{i\theta})d\theta \leq \int_0^{2\pi} \psi\circ f(e^{i\theta})d\theta < \int_0^{2\pi} u\circ f(e^{i\theta})d\theta + \epsilon.$$

This for any $\epsilon > 0$ hence $\int_0^{2\pi} u\circ g(re^{i\theta})d\theta \leq \int_0^{2\pi} u\circ f(e^{i\theta})d\theta$. □

Assume that f is proper then our Theorem 1 generalizes Littlewood's Subordination theorem to the situations where g is not subordinate to f. So on taking $u = |.|^p$, $\infty > p > 0$ we get

$$\int_0^{2\pi} |g(e^{i\theta})|^p d\theta \leq \int_0^{2\pi} |f(e^{i\theta})|^p d\theta.$$

One has the following interesting consequence. If two proper holomorphic functions g and f are such that $g(0) = f(0)$ and $g(\mathbb{D}) = f(\mathbb{D})$, then they have the same hardy norm. From now we say that $g \in \mathscr{O}(\mathbb{D},\mathbb{C})$ is subordinate to $f \in \mathscr{O}(\mathbb{D},\mathbb{C})$ if $g(0) = f(0)$ and $g(\mathbb{D}) \subset f(\mathbb{D})$.The following theorem gives a relation between the zero sets of subordinate and proper superordinate functions.

Theorem 3. *Let $f \in \mathscr{O}(\mathbb{D},\mathbb{C}) \cap C(\overline{\mathbb{D}},\mathbb{C})$ proper. Then for all $g \in \mathscr{O}(\mathbb{D},\mathbb{C})$ such that $g(\mathbb{D}) \subset f(\mathbb{D})$ and $g(0) = f(0)$ we have*

$$\sum_{b\in\mathbb{D}\cap g^{-1}(0)} m_b \log|b| \geq \sum_{a\in f^{-1}(0)} m_a \log|a|,$$

moreover if g is proper and $g(\mathbb{D}) = f(\mathbb{D})$ then we get

$$\sum_{b\in g^{-1}(0)} m_b \log|b| = \sum_{a\in f^{-1}(0)} m_a \log|a|.$$

With the convention $\sum_{a\in f^{-1}(0)} m_a \log|a| = 0$ if f doesn't vanish.

Proof. If $g(0) = f(0) = 0$. Then we get $-\infty = \sum_{b\in\mathbb{D}\cap g^{-1}(0)} m_b \log|b| = \sum_{a\in f^{-1}(0)} m_a \log|a|$. If $\mathbb{T}\cap f^{-1}(0)$ is not empty, then all the zeros are on $\mathbb{T}$ hence

$$0 = \sum_{b\in\mathbb{D}\cap g^{-1}(0)} m_b \log|b| = \sum_{a\in f^{-1}(0)} m_a \log|a|.$$

Assume that $f(0) = g(0) \neq 0$ and $\mathbb{T}\cap f^{-1}(0)$ is empty, then by Jesen's formula we have

$$\log|f(0)| + \sum_{a\in f^{-1}(0)} m_a \log\frac{1}{|a|} = \int_0^{2\pi} \log|f(e^{i\theta})|d\theta$$

$$\log|g(0)| + \sum_{b\in D_r\cap g^{-1}(0)} m_b \log\frac{r}{|b|} = \int_0^{2\pi} \log|g(re^{i\theta})|d\theta, \text{ for } 0 \leq r < 1.$$

By Theorem 1 we get

$$\log|g(0)| + \sum_{b\in D_r\cap g^{-1}(0)} m_b \log\frac{r}{|b|} \leq \log|f(0)| + \sum_{a\in f^{-1}(0)} m_a \log\frac{1}{|a|}.$$

As $g(0) = f(0) \neq 0$ then for all $r \in]0,1[$ we have

$$\sum_{b\in D_r\cap g^{-1}(0)} m_b \log\frac{r}{|b|} \leq \sum_{a\in f^{-1}(0)} m_a \log\frac{1}{|a|}$$

hence

$$\sum_{b\in\mathbb{D}\cap g^{-1}(0)} m_b \log\frac{1}{|b|} \leq \sum_{a\in f^{-1}(0)} m_a \log\frac{1}{|a|}.$$

□

It is well known for a continuous function $f:\mathbb{D}\to\mathbb{D}$ that, if $f(0)\neq 0$ then there is $r>0$ depending on f such that $f(z)\neq 0$ for $z\in D(0,r)$. Here we will prove that for holomorphic functions the choice of r may not depend of f.

Corollary 4. *Let $g\in\mathscr{O}(\mathbb{D},\mathbb{D})$ with $g(0)\neq 0$ then, g doesn't vanish in $D(0,|g(0)|)$.*

Proof. Set $a_0=g(0)$ and $f(z)=\frac{z+a_0}{1+\bar{a}_0 z}$ then f is proper and has a simple zero at $-a_0$. Notice that g is subordinate to f. Now assume that there is $b_0\in D(0,|g(0)|)$ such that $g(b_0)=0$, then

$$\sum_{a\in f^{-1}(0)} m_a \log|a| \geq \log|-a_0| > \log|b_0| \geq \sum_{b\in\mathbb{D}\cap g^{-1}(0)} m_b \log|b|.$$

This is in contradiction with the theorem above. □

For $x\in\mathbb{D}^*$ we denote by $\mathscr{O}(\mathbb{D},\mathbb{D},x)$ the set of all holomorphic functions $f\in\mathscr{O}(\mathbb{D},\mathbb{D})$ with $f(0)=x$, then $D(0,|x|)$ is the largest disk in which none element of $\mathscr{O}(\mathbb{D},\mathbb{D},x)$ vanishes. Notice that Corollary 4 can be seen as a consequence of Schwarz Lemma and Corollary 6 a generalization of Corollary 4.

Corollary 5. *Let $X\subset\mathbb{C}$ be open and $f\in\mathscr{O}(\mathbb{D},X)\cap C(\overline{\mathbb{D}},X)$ be proper. Then for all $g\in\mathscr{O}(\mathbb{D},X)$ such that $g(\mathbb{D})\subset f(\mathbb{D})$ and $g(0)=f(0)$ we have*

$$\sum_{b\in g^{-1}(p)} m_b \log|b| \geq \sum_{a\in f^{-1}(p)} m_a \log|a|, \text{ for } p\in X.$$

With the convention $\sum_{a\in f^{-1}(p)} m_a\log|a|=0$ if the function $f-p$ doesn't vanish.

The corollary above gives an idea about the location of the solutions of certain equations. For instance we have.

Corollary 6. *Let $g\in\mathscr{O}(\mathbb{D},\mathbb{D},x)$ with $x\in\mathbb{D}^*$ then, the function $g-a$ doesn't vanish in the disk $D(0,\frac{|a-x|}{|1-\bar{x}a|})$ for all $a\in\mathbb{D}\setminus\{x\}$.*

2.1. **Lelong functional in $\mathbb{C}$.** Consider $\alpha(x)=\sum_{j=1}^N m_j\chi_{\{p_j\}}(x)$, where p_j are points in X and m_j are positive, then we define a disc functional

$$H^L_\alpha:\mathscr{O}(\mathbb{D},X)\to\mathbb{R}\cup\{-\infty\},\ H^L_\alpha(f)=\sum_{a\in\mathbb{D}}\alpha(f(a))m_a\log|a|,$$

which is called the Lelong functional with respect to α. Its envelope is the following

$$EH^L_\alpha(x)=\inf\{H^L_\alpha(f), f\in\mathscr{O}(\mathbb{D},X), f(0)=x\}.$$

In [5] it is proven that EH^L_α is subharmonic and it coincides with the Green function of X with several poles at $p_1,\ldots,p_N$ of weights $m_1\ldots,m_N$. The theorem below states that the $infimum$ in the definition of EH^L_α is actually a $minimum$ in a special case.

Theorem 7. *Let X be open in $\mathbb{C}$. If there is $F \in \mathscr{O}(\mathbb{D}, \mathbb{C})$ proper such that $F(\mathbb{D}) = X$, then for all $x \in X$ there is $f \in \mathscr{O}(\mathbb{D}, X)$ with $f(0) = x$ such that*

$$EH_\alpha^L(x) = H_\alpha^L(f) = \sum_{z\in\mathbb{D}} \alpha(f(z)) m_z \log|z|.$$

Proof. Let $x \in X$, ψ be an automorphism of $\mathbb{D}$ such that $F\circ\psi(0) = x$ and set $f = F\circ\psi$. For all $g \in \mathscr{O}(\mathbb{D}, X)$ with $g(0) = x$ we have

$$H_\alpha^L(f) = \sum_{a\in\mathbb{D}} \alpha(f(a)) m_a \log|a| = \sum_{p\in X} \alpha(p) \sum_{a\in f^{-1}(p)} m_a \log|a|.$$

$$H_\alpha^L(g) = \sum_{b\in\mathbb{D}} \alpha(g(b)) m_b \log|b| = \sum_{p\in X} \alpha(p) \sum_{b\in g^{-1}(p)} m_b \log|b|.$$

Then by Corollary 5 we get $H_\alpha^L(f) \le H_\alpha^L(g)$ hence

$$EH_\alpha^L(x) \le H_\alpha^L(f) \le \inf\{H_\alpha^L(g), g \in \mathscr{O}(\mathbb{D}, X), g(0) = x\} \le EH_\alpha^L(x).$$

□

Let $z_x \in F^{-1}(x)$ and set $\psi_x(z) = \frac{z+z_x}{1+\bar{z}_x z}$ on taking $\psi_x(z) = a$ we get

$$EH_\alpha^L(x) = \sum_{a\in\mathbb{D}} \alpha(F(a)) m_a \log\left|\frac{a - z_x}{1 - \bar{z}_x a}\right|.$$

Remark that the value of $EH_\alpha^L(x)$ doesn't depend on the choice of $z_x \in F^{-1}(x)$, $x \in X$.

2.1.1. *Lelong functional in $\mathbb{C}^n$.* Let $X \subset \mathbb{C}^n$ be open, consider $\alpha(x) = m\chi_{\{p\}}(x)$, where p is a point in X and m is nonnegative.

Theorem 8. *Assume that $X = X_1 \times \cdots \times X_n$ and $X_i \subset \mathbb{C}$, $i = 1, ..., n$ are jordan domains, then at each point $x \in X$ there is an extremal disc for EH_α^L in other word there is $f \in \mathscr{O}(\mathbb{D}, X)$ with $f(0) = x$ such that*

$$EH_\alpha^L(x) = H_\alpha^L(f) = \sum_{a\in\mathbb{D}} \alpha(f(a)) m_a \log|a|.$$

Proof. Denote F_i the Riemann mapping from $\mathbb{D}$ onto X_i, for $x \in X$ we consider ψ_i an automorphism of $\mathbb{D}$ such that $F_i \circ \psi_i(0) = x_i$. Take $z_i \in \mathbb{D}$ such that $F_i \circ \psi_i(z_i) = p_i$. Take $a \in \{z_1, \ldots, z_n\}$ such that $|a| = \max\{|z_i|, i = 1, \ldots, n\}$. We obtain an analytic disc f in X centered at x containing p on setting $f(z) = (F_1 \circ \psi_1(\frac{z_1}{a} z), \ldots, F_n \circ \psi_n(\frac{z_n}{a} z))$. We may assume that $a = z_1$. Let $g \in \mathscr{O}(\mathbb{D}, X)$ with $g(0) = x$ remark that

$$\sum_{a\in g^{-1}(p)} m_a \log|a| \ge \sum_{a\in g_1^{-1}(p_1)} m_a \log|a| \ge \sum_{a\in f_1^{-1}(p_1)} m_a \log|a| = \sum_{a\in f^{-1}(p)} \log|a|.$$

Hence $H_\alpha^L(g) \ge H_\alpha^L(f)$ for all $g \in \mathscr{O}(\mathbb{D}, X, x)$ then $EH_\alpha^L(x) = H_\alpha^L(f) = \alpha(f(a)) \log|a|$ where $f(a) = p$. □

Remark that on setting $z_i = F^{-1}(p_i)$ we can write EH_α^L as follow

$$EH_\alpha^L(x) = \alpha(p) \max\left\{\log\left|\frac{F^{-1}(x_i) - z_i}{1 - \bar{z}_i F^{-1}(x_i)}\right|, i = 1, \ldots, n\right\}.$$

Corollary 9. *Let g and f be two closed proper analytic discs in X, where $X \subset \mathbb{C}$ is open. Assume that $g(0) = f(0)$ and $g(\overline{\mathbb{D}}) = f(\overline{\mathbb{D}})$, then for any $u \in SH(X)$ bounded one has:*

$$\int_0^{2\pi} u \circ g(e^{i\theta})d\theta = \int_0^{2\pi} u \circ f(e^{i\theta})d\theta.$$

2.2. **Coefficient Inequalities.** If $g(z) = \sum b_n z^n$ is subordinate to a proper function $f(z) = \sum a_n z^n$, then the coefficients of f dominate those of g in a certain average sense.

Corollary 10. *Let $f(z) = \sum_{n=1}^{\infty} a_n z^n$ be proper analytic in $\mathbb{D}$, continuous on $\overline{\mathbb{D}}$ and $g(z) = \sum_{n=1}^{\infty} b_n z^n$ be analytic in $\mathbb{D}$. Suppose $g(\overline{\mathbb{D}}) \subset f(\overline{\mathbb{D}})$ and $b_1 \neq 0$. Then there exist $N > 2$ such that*

$$\sum_{k=2}^{n} |b_k|^2 \leq \sum_{k=1}^{n} |a_k|^2, \text{ for all } n = N+1, \ldots.$$

Proof. Set $A_n = \sum_{k=1}^{n} |a_k|^2$ and $B_n = \sum_{k=1}^{n} |b_k|^2$. By Parseval's relation,

$$\frac{1}{2\pi}\int_0^{2\pi} |f(e^{i\theta})|^2 d\theta = \sum_{k=1}^{\infty} |a_k|^2 = \lim_{n\to\infty} A_n.$$

So there is $N > 2$ such that $\lim_{n\to\infty} A_n \leq A_n + |b_1|^2$ for $n > N$. Then by Theorem 1,

$$\sum_{k=1}^{n} |b_k|^2 \leq \frac{1}{2\pi}\int_0^{2\pi} |g(e^{i\theta})|^2 d\theta \leq \frac{1}{2\pi}\int_0^{2\pi} |f(e^{i\theta})|^2 d\theta \leq \sum_{k=1}^{n} |a_k|^2 + |b_1|^2, \text{ for } n > N.$$

Hence

$$\sum_{k=2}^{n} |b_k|^2 \leq \sum_{k=1}^{n} |a_k|^2, \text{ for } n > N.$$

□

Corollary 11. *If $a_n = O(1)$, then $b_n = O(\sqrt{n})$ as $n \to \infty$.*

Here we compare the area of $f(D_r)$ with the area of $g(D_{r^2})$ for r small.

Corollary 12. *Let $f(z) = \sum_{n=1}^{\infty} a_n z^n$ be proper analytic in D_r, $g(z) = \sum_{n=1}^{\infty} b_n z^n$ be analytic in D_r and suppose $g(D_r) \subset f(D_r)$. Then*

$$\int\int_{|z|\leq r^2} |g'(\rho e^{i\theta})|^2 \rho d\rho d\theta \leq \int\int_{|z|\leq r} |f'(\rho e^{i\theta})|^2 \rho d\rho d\theta, \ 0 \leq r \leq \left(\frac{1}{e}\right)^{\frac{1}{2e}}.$$

Proof. The integrals represent the areas of the image (the multisheeted image) of D_{r^2} under g and the image of D_r under f. By Theorem 1 we have

$$\sum_{n=1}^{\infty} |b_n|^2 r^{2n} \leq \sum_{n=1}^{\infty} |a_n|^2 r^{2n}.$$

Notice that for $r \leq (\frac{1}{e})^{\frac{1}{2e}}$, the function $x \to xr^{2x}$ is less than 1 for all $x \geq 1$. Therefore $nr^{2n} \leq 1$ for all $n > 1$, then

$$\sum_{n=1}^{\infty} nr^{2n}|b_n|^2 r^{2n} \leq \sum_{n=1}^{\infty} |b_n|^2 r^{2n} \leq \sum_{n=1}^{\infty} |a_n|^2 r^{2n} \leq \sum_{n=1}^{\infty} n|a_n|^2 r^{2n}.$$

Hence

$$\sum_{n=1}^{\infty} n|b_n|^2 r^{2^{2n}} \leq \sum_{n=1}^{\infty} n|a_n|^2 r^{2n}.$$

This last inequality is equivalent to the inequality in the corollary. □

We can also compare the Hardy norm of the derivatives on D_r, $r \leq (\frac{1}{e})^{\frac{1}{e}}$. The proof of the corollary below is similar to that of the corollary above.

Corollary 13. *Let $f(z) = \sum_{n=1}^{\infty} a_n z^n$ be proper analytic in D_r, $g(z) = \sum_{n=1}^{\infty} b_n z^n$ be analytic in D_r and suppose that $g(D_r) \subset f(D_r)$. Then*

$$\int_0^{2\pi} |g'(r^2 e^{i\theta})|^2 d\theta \leq \int_0^{2\pi} |f'(re^{i\theta})|^2 d\theta,\ 0 \leq r \leq \left(\frac{1}{e}\right)^{\frac{1}{e}}.$$

Acknowledgement. The second author thanks his supervisor Professor Armen Edigarian for his help to accomplish this work.

References

[1] P. L. Duren; univalent functions.
[2] Jan Wiegerinck; The Pluripolar hull of $\{w = e^{-1/z}\}$. Arkiv for Mat 38 2000 201-208.
[3] Thomas Ransford; Potential Theory in the Complex Plane.
[4] Maciej Klimek; Pluripotential Theory.
[5] F. Larusson and R. Sigurdson; Plurisubharmonic extremal function, Lelong numbers and Coherent Ideal Sheaves.
[6] Isabelle Chalendar; Fonctions Harmoniques, Classe de Nevanlinna, Espaces de Hardy, et une introduction aux operateurs de Hankel.

Science Park 904, Amsterdam
E-mail address: J.J.O.O.Wiegerinck@uva.nl

Jagiellonian University, Department of Mathematics
E-mail address: Ibrahim.Djire@im.uj.edu.pl